

# Talking with your doctor about your PI treatment plan?

Patient Name: \_\_\_\_\_

Date: \_\_\_\_\_

If you're living with PI, it's important to keep your doctor up to date about your condition. But talking during visits isn't always easy. Answering the questions below will help prepare you for your next appointment.\*



## How well is your PI being controlled?

\*Questions based on 2018 and 2023 National Treatment Survey, Immune Deficiency Foundation.

- 1 In the past 12 months, have you experienced recurrent lower respiratory tract infections that have resulted in bronchitis or pneumonia?

If YES, about how many times? \_\_\_\_\_

YES NO NOT SURE

1 2 3 MORE
- 2 In the past 12 months, have you experienced upper respiratory infections, such as sinusitis?

YES NO NOT SURE
- 3 Have you ever been told that you have bronchiectasis, asthma, COPD, or some other type of chronic lung condition?

YES NO NOT SURE
- 4 In the past 12 months, have you been prescribed antibiotics, antivirals, steroids, or other medications to treat, prevent, or manage a respiratory tract infection?

If YES, about how many times in the past 12 months? \_\_\_\_\_

YES NO

1 2 3 MORE
- 5 In the past 12 months, did you miss any days of school, work, or routine activities because you were sick?

If YES, about how many days did you miss? \_\_\_\_\_

YES NO

1-3 4 OR MORE
- 6 In the past 12 months, have you had an ER visit or been hospitalized due to a respiratory tract infection?

If YES, about how many times in the past 12 months? \_\_\_\_\_

YES NO

1 2 3 MORE
- 7 In the past 12 months, have you seen or been referred to a pulmonary specialist?

If YES, about how many times in the past 12 months? \_\_\_\_\_

YES NO

1 2 3 MORE
- 8 Has your doctor ever adjusted the dose or schedule of your Ig replacement therapy?

YES NO NOT SURE
- 9 Has your doctor ever switched your Ig replacement therapy?

YES NO NOT SURE

**If you answered YES to any question, you may be able to improve your PI protection.**

Consider showing your answers to your doctor. Ask if it might be time for a change in your Ig treatment.

## Is your PI under control?

Not all Ig treatments are the same. What works for one person may not work for someone else.


When people with PI continue to suffer from respiratory tract infections it is not uncommon for their healthcare provider to recommend changing to a different Iq.



EVERY PI PATIENT'S  
EXPERIENCE IS  
DIFFERENT, AND  
**YOUR JOURNEY  
IS UNIQUE**

## Changing Ig treatments is not unusual

A survey by the Immune Deficiency Foundation revealed that as many as **1 in 7 people** with PI switched from one Ig treatment to another in hopes of improving treatment results.\*



## Finding the right treatment can positively impact your overall wellbeing

\*2018 National Treatment Survey, Immune Deficiency Foundation.

Please see Important Safety Information on pages 5-7, and visit [asceniv.com](https://www.asceniv.com) for Full Prescribing Information, including complete BOXED WARNING.



DISCOVER THE  
ASCENIV™ DIFFERENCE

ASCENIV is an intravenous immune globulin (IVIG) for adults and adolescents (12-17 years of age) with PI.

**ASCENIV provides proven protection against infection and has a demonstrated safety profile.**

## Select Donor Profile

The only available IVIG product made with a proprietary method for donor screening AND plasma pooling

## Quality in Every Vial

Every vial includes a broad spectrum of antibodies, taken from plasma pooled from at least 1,000 donors

**ASCENIV is the only IVIG that uses ADMA's patented methodologies for donor screening and plasma pooling.**

ASCENIV meets FDA IVIG criteria, but also includes plasma from donors selected for their elevated antibody levels for certain respiratory pathogens

## Notes

[illegible]

Please see Important Safety Information on pages 5-7, and visit [asceniv.com](https://www.asceniv.com) for Full Prescribing Information, including complete BOXED WARNING.

SAVINGS AND SUPPORT MADE SIMPLE



ADMA ADvantage Ig is a dedicated patient support hub created for eligible patients who have been prescribed ASCENIV.

Ask your doctor today about enrolling!

Learn more about the ADvantage Ig Patient Support Program at [www.ASCENIV.com](http://www.ASCENIV.com)



DESIGNED TO DELIVER™

[www.asceniv.com](http://www.asceniv.com)

Please see Important Safety Information on pages 5-7, and visit [asceniv.com](http://asceniv.com) for Full Prescribing Information, including complete BOXED WARNING.

IMPORTANT SAFETY INFORMATION FOR ASCENIV™

**WARNING: RISK OF BLOOD CLOTS (THROMBOSIS), POOR KIDNEY FUNCTION, AND INABILITY TO FILTER WASTE FROM KIDNEYS. BLOOD CLOTS MAY OCCUR WITH INTRAVENOUS IMMUNE GLOBULIN PRODUCTS, INCLUDING ASCENIV.**

Before taking ASCENIV, talk to your doctor if you:

- Are of advanced age
- Are unusually sedentary (long periods of sitting down or inactive)
- Are taking estrogen-containing medicines (birth control pills, hormone replacement therapy)
- Have a permanent intravenous (IV) catheter
- Have hyperviscosity of the blood (diseases such as multiple myeloma or other causes of elevated proteins in the blood)
- Have cardiovascular (heart) problems or previous history of stroke

Thrombosis may occur even if you do not have any risk factors.

Serious kidney problems and death can also happen in certain patients who receive such products.

If you are at high risk of thrombosis or kidney problems, your doctor should adjust the dose of ASCENIV and will monitor you for signs and symptoms of thrombosis and viscosity, as well as kidney function.

What is ASCENIV (immune globulin intravenous, human–slra)?

ASCENIV (immune globulin intravenous, human–slra) is a prescription medicine to help adults and adolescents (12 to 17 years old) with primary immunodeficiency fight and prevent infections.

ASCENIV is for intravenous administration only. ASCENIV is made from healthy human blood/plasma.

Who should not use ASCENIV?

ASCENIV should not be used if you had a severe allergic reaction to human immune globulin or if you have been told by a doctor that you are immunoglobulin A (IgA)-deficient and have developed antibodies to IgA and hypersensitivity after exposure to a previous plasma product.

IMPORTANT SAFETY INFORMATION (cont’d)

What are possible warnings and precautions with taking ASCENIV™?

**Hypersensitivity.** Severe allergic reactions may occur with immune globulin products, including ASCENIV. If you have a severe allergic reaction, stop the infusion immediately and get medical attention. ASCENIV contains IgA. If you have known antibodies to IgA, you may have a greater risk of developing potentially severe allergic reactions.

If you take ASCENIV or a similar immune globulin product, you could experience a serious and life-threatening blood clot (thromboembolism). This may include pain and/or swelling of an arm or leg with warmth over the affected area, discoloration of an arm or leg, unexplained shortness of breath, chest pain or discomfort that worsens on deep breathing, unexplained rapid pulse, numbness, or weakness on one side of the body. If you are at risk, your doctor may decide to adjust the dose of ASCENIV. Your doctor will monitor you for any signs or symptoms of blood clots or poor blood flow in your arteries.

**Always tell your doctor immediately if your medical history is similar to what is described here, and especially if you experience any of these symptoms while taking ASCENIV.**

**Kidney problems or failure.** Kidney problems, kidney failure, and death may occur with use of human immune globulin products, especially those containing sucrose (sugar). ASCENIV does not contain sucrose.

If you have kidney disease or diseases with kidney involvement, your doctor should perform a blood test to assess your hydration level and kidney function before beginning immune globulin treatment and at appropriate intervals thereafter. If your doctor determines that kidney function is worsening, they may discontinue treatment. If your doctor determines you to be at risk, they may start your dose of ASCENIV at a safe level.

**People taking human immune globulin products, including ASCENIV, may experience hyperproteinemia (high levels of protein in the blood), hyponatremia (low levels of sodium in the blood), and hyperviscosity (poor blood flow). Your doctor may perform certain blood tests and monitor you to minimize any of the above risks.**

**Aseptic meningitis syndrome (AMS).** Aseptic meningitis is a non-infectious inflammation of the membranes that cover the brain. It causes a severe headache, which may occur with human immune globulin treatment, including ASCENIV. AMS usually happens within a few hours to 2 days after treatment. AMS is more commonly associated with higher doses of treatment and/or after rapid infusion. Your doctor may perform a neurological exam, including spinal tap (sampling fluid which surrounds the spinal cord) to evaluate your condition and to rule out other causes of meningitis.

**Hemolysis.** Hemolysis refers to the destruction of red blood cells. Immune globulin products, including ASCENIV, may contain certain antibodies that can result in the rupturing of red blood cells. Your doctor should monitor you for signs and symptoms of hemolysis, which may include additional confirmation tests.

Taking intravenous human immune globulin products may cause a build up of fluid in the lungs (pulmonary edema) that is unrelated to heart problems. Your doctor should monitor you for lung-related side effects and may conduct appropriate tests that can detect the presence of certain white blood cells (anti-neutrophil antibodies) in the drug or your blood. If needed, your doctor may decide to use oxygen or other respiratory methods to help your breathing.

IMPORTANT SAFETY INFORMATION (cont’d)

**Transmissible infectious agents.** Because ASCENIV™ is made from human blood, it may carry a risk of transmitting infectious agents such as viruses, the variant Creutzfeldt-Jakob disease (vCJD) agent, and, theoretically, the Creutzfeldt-Jakob disease (CJD) agent. Your doctor will report to the manufacturer any cases of suspected infections spread by the product.

**Interference with lab tests.** Because ASCENIV contains a variety of antibodies that are infused into your body, blood tests to determine antibody levels may provide misleading interpretations. Be sure to always tell your doctor, nurse, or lab technician of any medicines you are taking and that you are using ASCENIV.

**Interactions with medicines.** ASCENIV can make vaccines (like measles, mumps, rubella, and chicken pox vaccines) less effective in your body. Before you get any vaccines, tell your healthcare provider that you take ASCENIV.

What are other possible side effects of ASCENIV?

In clinical studies of ASCENIV, some patients experienced the following:

- Headache
- Sinus inflammation (sinusitis)
- Diarrhea
- Intestinal lining inflammation caused by virus (gastroenteritis)
- Common cold (nasopharyngitis)
- Upper respiratory tract infection
- Bronchitis
- Nausea

**These are not all the possible side effects of ASCENIV.**

**Talk to your healthcare provider about any side effect that bothers you or that does not go away.**

**You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088.**



Scan code to learn more



at **asceniv.com**



© 2025 ADMA Biologics. All rights reserved.  
ASCENIV™, DESIGNED TO DELIVER™, and ADvantage Ig™ are  
trademarks of ADMA Biologics.  
11180-22-IGG-03192025\_R00  
Visit [admabiologics.com](https://admabiologics.com)

**ASCENIV**™  
IMMUNE GLOBULIN INTRAVENOUS  
(HUMAN) — slra 10% LIQUID